

The study of $B_c^- \rightarrow X(3872)\pi^-(K^-)$ Decays in the Covariant Light-Front Approach

Wei Wang, Yue-Long Shen and Cai-Dian Lü

Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, P.R. China

Abstract

In the covariant light-front quark model, we calculate the form factors of $B_c^- \rightarrow J/\psi$ and $B_c^- \rightarrow X(3872)$. Since the factorization of the exclusive processes $B_c^- \rightarrow J/\psi\pi^-(K^-)$ and $B_c^- \rightarrow X(3872)\pi^-(K^-)$ can be proved in the soft-collinear effective theory, we can get the branching ratios for these decays easily from the form factors. Taking the uncertainties into account, our results for the branching ratio of $B_c^- \rightarrow J/\psi\pi^-(K^-)$ are consistent with the previous studies. By identifying $X(3872)$ as a 1^{++} charmonium state, we obtain $\mathcal{BR}(B_c^- \rightarrow X(3872)\pi^-) = (1.7_{-0.6}^{+0.7+0.1+0.4}) \times 10^{-4}$ and $\mathcal{BR}(B_c^- \rightarrow X(3872)K^-) = (1.3_{-0.5}^{+0.5+0.1+0.3}) \times 10^{-5}$. If assuming $X(3872)$ as a 1^{--} state, the branching ratios will be one order magnitude larger than those of 1^{++} state. These results can be easily used to test the charmonium description for this mysterious meson $X(3872)$ at LHCb experiment.

I. INTRODUCTION

$X(3872)$ was first observed by Belle in the exclusive decay $B^\pm \rightarrow K^\pm X \rightarrow K^\pm \pi^+ \pi^- J/\psi$ [1], and subsequently confirmed by CDF, D0 and BaBar collaborations in various decay and production channels [2]. At present a definite answer on its internal properties is not well established, but the current experimental data strongly support a 1^{++} state [3]. Enormous interest in $\bar{c}c$ resonance spectroscopy study followed this discovery and there exists many interpretations for this meson. The first and most natural assignment of this state is the first radial excitation of $1P$ charmonium state χ_{c1} [4]. However, this interpretation has encountered two difficulties: its decay width (< 2.3 MeV, 95% C.L.) is tiny compared with other charmonia; there is a gap of about 100 MeV between the measured mass and the quark model prediction [5]. Motivated by these two difficulties, many non-charmonium explanations were proposed, such as $\bar{c}cg$ hybrid meson [6], glueball [7], diquark cluster [8], and molecular state [9]. But in fact, there is few experimental data which could provide a clear discrimination among these descriptions and this makes the situation more obscure. Recently the CLQCD collaboration studied the mass for the first excited states of 1^{++} charmonium and found it is consistent with the measured mass of $X(3872)$ [10]. The consistence indicates that $X(3872)$ can be the first radial excited state of χ_{c1} and it seems that the mass difficulty trails off. Now, in order to investigate the structure of this meson more clearly, a large amount of experimental data and theoretical studies on the productions and decays of $X(3872)$ are strongly deserved.

In $B_{u,d,s}$ decays involving charmonium final states, the emitted meson is a heavy charmonium. The non-factorizable contribution should be large to induce large uncertainties [11]. As the energy release is limited, these decays may also be polluted by the final state interactions which are non-perturbative in nature. But fortunately the production of charmonia in B_c decays could provide a unique insight to these mesons. Since the emitted meson here is a light meson (π or K), the factorization of $B_c \rightarrow (\bar{c}c)M$ (M is a light meson) could be proved in the framework of Soft-Collinear-Effective-Theory (SCET) to all orders of the strong coupling constant in heavy quark limit which is similar with $\bar{B}^0 \rightarrow D^+ \pi^-$ and $B^- \rightarrow D^0 \pi^-$ [12]. The decay matrix element can be decomposed into the $B_c \rightarrow (\bar{c}c)$ form factor and a convolution of a short distance coefficient with the light-cone wave function of the emitted light meson.

Although SCET provides a powerful framework to study the factorization of the exclusive modes, the non-perturbative form factors could not be directly studied. We can only extract them via the experimental data or rely on some non-perturbative method. In the present paper, we will use the light-front quark model to calculate these $B_c \rightarrow M(\bar{c}c)$ form factors. As pointed out in [13], the light front QCD approach has some unique features which are particularly suitable to describe a hadronic bound state. The light-front quark model [14, 15, 16] can provide a relativistic treatment of the movement of the hadron and also give a fully treatment of the hadron spin by using the so-called Melosh rotation. Light-

front wave functions, which describe the hadron in terms of their fundamental quark and gluon degrees of freedom, are independent of the hadron momentum and thus are explicitly Lorentz invariant. Furthermore, in covariant light-front approach[17], the spurious contribution which is dependent on the orientation of the light-front is eliminated by including the zero-mode contributions properly. This covariant model has been successfully extended to study the decay constants and form factors of the ground-state s -wave and low-lying p -wave mesons [18, 19, 20].

The paper is organized as follows. The formalism for the form factor calculations, taking $B_c^- \rightarrow J/\psi$ as an example, is presented in the next section. The numerical results for form factors and decay rates of $B_c^- \rightarrow J/\psi\pi^-(K^-)$, $B_c^- \rightarrow J/\psi\rho^-(K^{*-})$, $B_c^- \rightarrow X(3872)\pi^-(K^-)$ and $B_c^- \rightarrow X(3872)\rho^-(K^{*-})$ are given in Section III. The conclusion is given in Section IV.

II. CALCULATION OF THE FORM FACTORS AND THE BRANCHING RATIOS

In the following, we use X to denote $X(3872)$ for simplicity. Different with $B_{u,d,s}$ mesons, the B_c^- system consists of two heavy quarks b and c , which can decay individually. Here we will consider b decays while $\bar{c}$ acts as a spectator. At the quark level, $B_c^- \rightarrow J/\psi\pi^-$ and $B_c^- \rightarrow X(3872)\pi^-$ decays are characterized by $b \rightarrow (cd\bar{u})$ transition and the corresponding effective Hamiltonian is given by [21] :

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* [C_1(\mu) O_1(\mu) + C_2(\mu) O_2(\mu)] + \text{H.c.}, \quad (1)$$

where V_{ij} are the corresponding CKM matrix elements. The local four-quark operators $O_{1,2}$ are defined by:

$$O_1(\mu) = (\bar{c}_\alpha b_\beta)_{V-A} (\bar{d}_\beta u_\alpha)_{V-A}, \quad O_2(\mu) = (\bar{c}_\alpha b_\alpha)_{V-A} (\bar{d}_\beta u_\beta)_{V-A}, \quad (2)$$

where α and β are the color indices. Since the four quarks in the operators are different with each other, there is no penguin contribution and thus there is no CP violation. The left handed current is defined as $(\bar{q}_\alpha q_\beta)_{V-A} = \bar{q}_\alpha \gamma_\nu (1 - \gamma_5) q_\beta$. For $b \rightarrow (cs\bar{u})$ transition, V_{ud}^* is replaced by V_{us}^* while d quark field in the four-quark operator is replaced by s . With the effective Hamiltonian given above, the matrix element for the $B_c^- \rightarrow J/\psi\pi^-$ transition can be expressed as:

$$\mathcal{M} = \langle J/\psi(P'', \varepsilon''^*) \pi | \mathcal{H}_{eff} | B_c^-(P') \rangle = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* a_1(\mu) \langle J/\psi(P'', \varepsilon''^*) \pi | O_2(\mu) | B_c^-(P') \rangle, \quad (3)$$

with $P'^{(\prime\prime)}$ being the incoming (outgoing) momentum, ε''^* the polarization vector of J/ψ and $a_1 = C_2 + C_1/3$ the Wilson coefficient.

In the effective Hamiltonian, the degrees of freedom heavier than b quark mass m_b scale are included in the Wilson coefficients which can be calculated using the perturbation theory.

Then the left task is to calculate the operators' matrix elements between the B_c^- meson state and the final states, which suffers large uncertainties. Nevertheless, the problem becomes tractable if factorization becomes applicable. Thanks to the development of SCET, the proof of the factorization can be accomplished in an elegant way [22, 23]. In SCET, the heavy meson is described by the heavy quarks h_v and soft gluons A_s in its rest frame; the final state light meson moves very fast and it is described by the collinear quarks ξ_c and collinear gluons A_c . In Ref. [12], it has been shown that the collinear gluons do not connect to the particle in the heavy meson while the soft gluons don't connect to those in the light meson to all orders in α_s and leading power in Λ_{QCD}/m_{B_c} . In a phenomenal language, the non-factorizable diagrams cancel with each other because of color transparency. Furthermore, there is no annihilation contribution as the quarks in the final state meson are different with each other. Thus the decay amplitude can be expressed as the product of $B_c \rightarrow J/\psi$ form factor and a convolution of a short distance Wilson coefficient with non-perturbative light-cone distribution amplitude of the light meson. Without the higher order QCD corrections, the convolution is reduced to the decay constant of the light meson.

The form factors for $B_c \rightarrow J/\psi$ and $B_c \rightarrow X(3872)$ (1^{++} state) transitions induced by the vector and axial-vector currents are defined by:

$$\langle J/\psi(P'', \varepsilon''^*) | V_\mu | B_c^-(P') \rangle = -\frac{1}{m_{B_c} + m_{J/\psi}} \epsilon_{\mu\nu\alpha\beta} \varepsilon''^{*\nu} P^\alpha q^\beta V^{PV}(q^2), \quad (4)$$

$$\begin{aligned} \langle J/\psi(P'', \varepsilon''^*) | A_\mu | B_c^-(P') \rangle = & i \left\{ (m_{B_c} + m_{J/\psi}) \varepsilon''^* A_1^{PV}(q^2) - \frac{\varepsilon''^* \cdot P}{m_{B_c} + m_{J/\psi}} P_\mu A_2^{PV}(q^2) \right. \\ & \left. - 2m_{J/\psi} \frac{\varepsilon''^* \cdot P}{q^2} q_\mu [A_3^{PV}(q^2) - A_0^{PV}(q^2)] \right\}, \end{aligned} \quad (5)$$

$$\begin{aligned} \langle X(P'', \varepsilon'') | V_\mu | B_c^-(P') \rangle = & (m_{B_c} - m_X) \varepsilon''^* V_1^{PA}(q^2) - \frac{\varepsilon''^* \cdot P'}{m_{B_c} - m_X} P'_\mu V_2^{PA}(q^2) \\ & - 2m_X \frac{\varepsilon''^* \cdot P'}{q^2} q_\mu [V_3^{PA}(q^2) - V_0^{PA}(q^2)], \end{aligned} \quad (6)$$

$$\langle X(P'', \varepsilon'') | A_\mu | B_c^-(P') \rangle = -\frac{i}{m_{B_c} - m_X} \epsilon_{\mu\nu\rho\sigma} \varepsilon''^{\nu} P^\rho q^\sigma A^{PA}(q^2). \quad (7)$$

where $P = P' + P''$, $q = P' - P''$ and the convention $\epsilon_{0123} = 1$ is adopted. To cancel the poles at $q^2 = 0$, we must have $A_3^{PV}(0) = A_0^{PV}(0)$, $V_3^{PA}(0) = V_0^{PA}(0)$. The form factor $A_3^{PV}(V_3^{PA})$ is related to other form factors by:

$$\begin{aligned} A_3^{PV}(q^2) &= \frac{m_{B_c} + m_{J/\psi}}{2m_{J/\psi}} A_1^{PV}(q^2) - \frac{m_{B_c} - m_{J/\psi}}{2m_{J/\psi}} A_2^{PV}(q^2), \\ V_3^{PA}(q^2) &= \frac{m_{B_c} - m_X}{2m_X} V_1^{PA}(q^2) - \frac{m_{B_c} + m_X}{2m_X} V_2^{PA}(q^2). \end{aligned} \quad (8)$$

Following the notation in Refs. [17, 18], we use the light-front decomposition of the momentum $P' = (P'^-, P'^+, P'_\perp)$, where $P'^\pm = P'^0 \pm P'^3$, so that $P'^2 = P'^+ P'^- - P'^2_\perp$ and work in the $q^+ = 0$ frame. The incoming and outgoing mesons have the momentum $P' = p'_1 + p_2$ and $P'' = p''_1 + p_2$, respectively. The quark and antiquark inside the incoming




FIG. 1: Feynman diagram for $B_c \rightarrow J/\psi(X(3872))$ decay amplitudes. The X in the diagram denotes the $V - A$ transition vertex while the meson-quark-antiquark vertices are given in the text.

(outgoing) meson have the mass $m_1^{(')}$ and m_2 whose momenta are denoted as $p_1^{(')}$ and p_2 respectively. These momenta can be expressed in terms of the internal variables $(x_i, p'_\perp)$ as:

$$p_{1,2}^+ = x_{1,2}P'^+, \quad p'_{1,2\perp} = x_{1,2}P'_\perp \pm p'_\perp, \quad (9)$$

with $x_1 + x_2 = 1$. Using these internal variables, we can define some other useful quantities for the incoming meson:

$$\begin{aligned} M_0'^2 &= (e'_1 + e'_2)^2 = \frac{p_\perp'^2 + m_1'^2}{x_1} + \frac{p_\perp'^2 + m_2^2}{x_2}, & \widetilde{M}_0' &= \sqrt{M_0'^2 - (m_1' - m_2)^2}, \\ e_i^{(')} &= \sqrt{m_i^{(')2} + p_\perp'^2 + p_z'^2}, & p_z' &= \frac{x_2 M_0'}{2} - \frac{m_2^2 + p_\perp'^2}{2x_2 M_0'}. \end{aligned} \quad (10)$$

$e_i^{(')}$ can be interpreted as the energy of the quark or the antiquark and M_0' can be viewed kinematic invariant mass of the meson system. To calculate the amplitude for the transition form factor, we need the following Feynman rules for the meson-quark-antiquark vertices $(i\Gamma'_M)^1$:

$$i\Gamma'_P = H'_P \gamma_5, \quad (11)$$

$$i\Gamma'_V = iH'_V [\gamma_\mu - \frac{1}{W'_V} (p'_1 - p_2)_\mu], \quad (12)$$

$$i\Gamma'_A = -H'_A [\gamma_\mu + \frac{1}{W'_A} (p'_1 - p_2)_\mu] \gamma_5. \quad (13)$$

Here and in the following, we use the subscript A to denote the axial-vector with the quantum numbers $J^{PC} = 1^{++}$. For the outgoing meson, we should use $i(\gamma_0 \Gamma_M^\dagger \gamma_0)$ for the corresponding vertices.

In the conventional light-front quark model, the constituent quarks are required to be on mass shell and the physical quantities can be extracted from the plus component of the corresponding current matrix elements. However, this framework suffers the problem of

¹ We use a different phase $-i$ with Ref. [18] for the incoming axial-vector vertex.

non-covariance and miss the zero-mode contributions. To solve this problem, Jaus proposed the covariant light-front approach which can deal with the zero mode contributions systematically [17]. Decay constants and form factors can be calculated in terms of Feynman momentum loop integrals which are manifestly covariant. In this framework, the lowest order contribution to a form factor is depicted in Fig. 1. For the $P \rightarrow V$ transition, the decay amplitudes are:

$$\mathcal{B}_\mu^{PV} = -i^3 \frac{N_c}{(2\pi)^4} \int d^4 p'_1 \frac{H'_P(iH''_V)}{N'_1 N''_1 N_2} S_{\mu\nu}^{PV} \varepsilon^{\prime\prime*\nu}, \quad (14)$$

where $N_1^{\prime(n)} = p_1^{\prime(n)2} - m_1^{\prime(n)2} + i\epsilon$, $N_2 = p_2^2 - m_2^2 + i\epsilon$ and

$$\begin{aligned} S_{\mu\nu}^{PV} &= (S_V^{PV} - S_A^{PV})_{\mu\nu} \\ &= \text{Tr} \left[\left(\gamma_\nu - \frac{1}{W_V''} (p_1'' - p_2)_\nu \right) (\not{p}_1'' + m_1'') (\gamma_\mu - \gamma_\mu \gamma_5) (\not{p}_1' + m_1') \gamma_5 (-\not{p}_2 + m_2) \right] \\ &= -2i\epsilon_{\mu\nu\alpha\beta} \left\{ p_1'^\alpha P^\beta (m_1'' - m_1') + p_1'^\alpha q^\beta (m_1'' + m_1' - 2m_2) + q^\alpha P^\beta m_1' \right\} \\ &\quad + \frac{1}{W_V''} (4p_{1\nu}' - 3q_\nu - P_\nu) i\epsilon_{\mu\alpha\beta\rho} p_1'^\alpha q^\beta P^\rho \\ &\quad + 2g_{\mu\nu} \left\{ m_2(q^2 - N_1' - N_1'' - m_1'^2 - m_1''^2) - m_1'(M''^2 - N_1'' - N_2 - m_1'^2 - m_2^2) \right. \\ &\quad \left. - m_1''(M'^2 - N_1' - N_2 - m_1'^2 - m_2^2) - 2m_1' m_1'' m_2 \right\} \\ &\quad + 8p_{1\mu}' p_{1\nu}' (m_2 - m_1') - 2(P_\mu q_\nu + q_\mu P_\nu + 2q_\mu q_\nu) m_1' + 2p_{1\mu}' P_\nu (m_1' - m_1'') \\ &\quad + 2p_{1\mu}' q_\nu (3m_1' - m_1'' - 2m_2) + 2P_\mu p_{1\nu}' (m_1' + m_1'') + 2q_\mu p_{1\nu}' (3m_1' + m_1'' - 2m_2) \\ &\quad + \frac{1}{2W_V''} (4p_{1\nu}' - 3q_\nu - P_\nu) \left\{ 2p_{1\mu}' [M'^2 + M''^2 - q^2 - 2N_2 + 2(m_1' - m_2)(m_1'' + m_2)] \right. \\ &\quad \left. + q_\mu [q^2 - 2M'^2 + N_1' - N_1'' + 2N_2 - (m_1' + m_1'')^2 + 2(m_1' - m_2)^2] \right. \\ &\quad \left. + P_\mu [q^2 - N_1' - N_1'' - (m_1' + m_1'')^2] \right\}. \end{aligned} \quad (15)$$

In practice, we use the light-front decomposition of the loop momentum and have to perform the integration over the minus component using the contour method, as the covariant vertex functions can not be determined by solving the bound state equation. If the covariant vertex functions are not singular when performing the integration, the transition amplitude will pick up the singularities in the antiquark propagator. The integration leads to:

$$\begin{aligned} N_1^{\prime(n)} &\rightarrow \hat{N}_1^{\prime(n)} = x_1 (M^{\prime(n)2} - M_0^{\prime(n)2}), \\ H_M^{\prime(n)} &\rightarrow h_M^{\prime(n)}, \\ W_M'' &\rightarrow w_M'', \\ \int \frac{d^4 p'_1}{N'_1 N''_1 N_2} H'_P H''_V S &\rightarrow -i\pi \int \frac{dx_2 d^2 p'_\perp}{x_2 \hat{N}_1' \hat{N}_1''} h'_P h''_V \hat{S}, \end{aligned} \quad (16)$$

where

$$M_0^{\prime(n)2} = \frac{p_\perp^{\prime(n)2} + m_1^{\prime(n)2}}{x_1} + \frac{p_\perp^{\prime(n)2} + m_2^2}{x_2}, \quad (17)$$

with $p'_\perp = p'_\perp - x_2 q_\perp$. The explicit forms of h'_M and w'_M for the pseudoscalar, vector and axial-vector 1^{++} are given by [18]

$$\begin{aligned} h'_P &= h'_V = (M'^2 - M_0'^2) \sqrt{\frac{x_1 x_2}{N_c}} \frac{1}{\sqrt{2\widetilde{M}_0'}} \varphi', \\ h'_A &= (M'^2 - M_0'^2) \sqrt{\frac{x_1 x_2}{N_c}} \frac{1}{\sqrt{2\widetilde{M}_0'}} \frac{\widetilde{M}_0'^2}{2\sqrt{2}M_0'} \varphi'_p, \end{aligned} \quad (18)$$

$$w'_V = M'_0 + m'_1 + m_2, \quad w'_A = \frac{\widetilde{M}_0'^2}{m'_1 - m_2}, \quad (19)$$

where φ' and φ'_p are the phenomenological light-front momentum distribution amplitudes for s -wave and p -wave mesons, respectively. After this integration, the conventional light-front model is recovered but manifestly the covariance is lost as it receives additional spurious contributions proportional to the lightlike four vector $\tilde{\omega} = (\tilde{\omega}^-, \tilde{\omega}^+, \tilde{\omega}_\perp) = (1, 0, 0_\perp)$. The spurious contributions can be eliminated by including the zero mode contribution which amounts to performing the p^- integration in a proper way [17, 18].

By using Eqs. (15)–(19) and the integration rules in Refs. [17, 18], one arrives at

$$\begin{aligned} g(q^2) &= -\frac{N_c}{16\pi^3} \int dx_2 d^2 p'_\perp \frac{2h'_P h''_V}{x_2 \hat{N}'_1 \hat{N}''_1} \left\{ x_2 m'_1 + x_1 m_2 + (m'_1 - m''_1) \frac{p'_\perp \cdot q_\perp}{q^2} + \frac{2}{w''_V} \left[p'^2_\perp + \frac{(p'_\perp \cdot q_\perp)^2}{q^2} \right] \right\}, \\ f(q^2) &= \frac{N_c}{16\pi^3} \int dx_2 d^2 p'_\perp \frac{h'_P h''_V}{x_2 \hat{N}'_1 \hat{N}''_1} \left\{ 2x_1 (m_2 - m'_1) (M_0'^2 + M_0''^2) - 4x_1 m''_1 M_0'^2 + 2x_2 m'_1 q \cdot P \right. \\ &\quad + 2m_2 q^2 - 2x_1 m_2 (M'^2 + M''^2) + 2(m'_1 - m_2) (m'_1 + m''_1)^2 + 8(m'_1 - m_2) \left[p'^2_\perp + \frac{(p'_\perp \cdot q_\perp)^2}{q^2} \right] \\ &\quad + 2(m'_1 + m''_1) (q^2 + q \cdot P) \frac{p'_\perp \cdot q_\perp}{q^2} - 4 \frac{q^2 p'^2_\perp + (p'_\perp \cdot q_\perp)^2}{q^2 w''_V} \left[2x_1 (M'^2 + M_0'^2) - q^2 - q \cdot P \right. \\ &\quad \left. \left. - 2(q^2 + q \cdot P) \frac{p'_\perp \cdot q_\perp}{q^2} - 2(m'_1 - m''_1) (m'_1 - m_2) \right] \right\}, \\ a_+(q^2) &= \frac{N_c}{16\pi^3} \int dx_2 d^2 p'_\perp \frac{2h'_P h''_V}{x_2 \hat{N}'_1 \hat{N}''_1} \left\{ (x_1 - x_2) (x_2 m'_1 + x_1 m_2) - [2x_1 m_2 + m''_1 + (x_2 - x_1) m'_1] \frac{p'_\perp \cdot q_\perp}{q^2} \right. \\ &\quad \left. - 2 \frac{x_2 q^2 + p'_\perp \cdot q_\perp}{x_2 q^2 w''_V} [p'_\perp \cdot p''_\perp + (x_1 m_2 + x_2 m'_1) (x_1 m_2 - x_2 m''_1)] \right\}, \end{aligned} \quad (20)$$

while the physical form factors are related to the above functions by

$$\begin{aligned} V^{PV}(q^2) &= -(m_{B_c} + m_{J/\psi}) g(q^2), \quad A_1^{PV}(q^2) = -\frac{f(q^2)}{m_{B_c} + m_{J/\psi}}, \\ A_2^{PV}(q^2) &= (m_{B_c} + m_{J/\psi}) a_+(q^2). \end{aligned} \quad (21)$$

The extension to $P \rightarrow A$ transitions is straightforward:

$$\mathcal{B}_\mu^{PA} = -i^3 \frac{N_c}{(2\pi)^4} \int d^4 p'_1 \frac{H'_P H''_A}{N'_1 N''_1 N_2} S_{\mu\nu}^{PA} \varepsilon^{\mu*\nu}, \quad (22)$$

where

$$\begin{aligned}
S_{\mu\nu}^{PA} &= (S_V^{PA} - S_A^{PA})_{\mu\nu} \\
&= \text{Tr} \left[\left(\gamma_\nu - \frac{1}{W_A''} (p_1'' - p_2)_\nu \right) \gamma_5 (\not{p}_1'' + m_1'') (\gamma_\mu - \gamma_\mu \gamma_5) (\not{p}_1' + m_1') \gamma_5 (-\not{p}_2 + m_2) \right] \\
&= \text{Tr} \left[\left(\gamma_\nu - \frac{1}{W_A''} (p_1'' - p_2)_\nu \right) (\not{p}_1'' - m_1'') (\gamma_\mu \gamma_5 - \gamma_\mu) (\not{p}_1' + m_1') \gamma_5 (-\not{p}_2 + m_2) \right]. \quad (23)
\end{aligned}$$

By comparing eq. (15) and eq. (23), we have $S_{V(A)}^{PA} = S_{A(V)}^{PV}$ with the replacement $m_1'' \rightarrow -m_1''$, $W_V'' \rightarrow W_A''$, except for a phase i . Consequently, the form factors of $B \rightarrow A$ can be related to the $B \rightarrow V$ form factors through:

$$\begin{aligned}
\ell^A(q^2) &= f(q^2) \text{ with } (m_1'' \rightarrow -m_1'', h_V'' \rightarrow h_A'', w_V'' \rightarrow w_A''), \\
q^A(q^2) &= g(q^2) \text{ with } (m_1'' \rightarrow -m_1'', h_V'' \rightarrow h_A'', w_V'' \rightarrow w_A''), \\
c_+^A(q^2) &= a_+(q^2) \text{ with } (m_1'' \rightarrow -m_1'', h_V'' \rightarrow h_A'', w_V'' \rightarrow w_A''), \quad (24)
\end{aligned}$$

where we should be cautious that the replacement of $m_1'' \rightarrow -m_1''$ can not be applied to m_1'' in w'' and h'' .

$$\begin{aligned}
A^{PA}(q^2) &= -(m_{B_c} - m_X) q(q^2), \quad V_1^{PA}(q^2) = -\frac{\ell(q^2)}{m_{B_c} - m_X}, \\
V_2^{PA}(q^2) &= (m_{B_c} - m_X) c_+(q^2). \quad (25)
\end{aligned}$$

In the above expressions for the form factors, there are many terms containing $(p'_\perp \cdot q_\perp)/q^2$ in the integrand. These terms can make non-trivial contributions together with $h_M''/\hat{N}_1''$. In the calculation, we make a Taylor expansion for $h_M''/\hat{N}_1''$ as:

$$\frac{h_V''}{\hat{N}_1''} = \frac{h_V''}{\hat{N}_1''} \Big|_{p_\perp'^2 \rightarrow p_\perp'^2} - 2 x_2 p'_\perp \cdot q_\perp \left(\frac{d}{dp_\perp'^2} \frac{h_V''}{\hat{N}_1''} \right)_{p_\perp'^2 \rightarrow p_\perp'^2} + \mathcal{O}(x_2^2 q^2). \quad (26)$$

Then terms containing $p'_\perp \cdot q_\perp$ can be simplified using the following equation:

$$\int d^2 p'_\perp \frac{(p'_\perp \cdot q_\perp)^2}{q^2} = -\frac{1}{2} \int d^2 p'_\perp p_\perp'^2. \quad (27)$$

Now it is straightforward to obtain the decay width:

$$\Gamma(B_c^- \rightarrow J/\psi \pi^-) = \frac{|G_F V_{cb} V_{ud}^* a_1 f_\pi m_{B_c}^2 A_0^{PV}(0)|^2}{32\pi m_{B_c}} (1 - r_{J/\psi}^2), \quad (28)$$

where $r_{J/\psi} = \frac{m_{J/\psi}}{m_{B_c}}$. For the decays involving K^- , the factor $V_{ud}^* f_\pi$ is replaced by $V_{us}^* f_K$; while for $B_c^- \rightarrow X \pi^- (K^-)$, $A_0^{PV}(0)(r_{J/\psi})$ is replaced by $V_0^{PA}(0)(r_X)$.

TABLE I: The input parameters m_q and β (in unit of GeV) in the Gaussian-type light-front wave function (29).

m_c	m_b	β_{B_c}	$\beta_{J/\psi}$	β_X
1.4	4.4	0.870 ± 0.100	$0.631^{+0.06}_{-0.04}$	0.720 ± 0.100

III. NUMERICAL RESULTS AND DISCUSSION

To perform numerical calculations we need to specify the input parameters in the covariant light-front framework. The $\bar{q}q$ meson state is described by the light-front wave function which can be obtained by solving the relativistic Schrödinger equation with a phenomenological potential. But in fact except for some special cases, the solution is not obtainable at present. We prefer to employ a phenomenological wave function to describe the hadronic structure. In the present work, we shall use the simple Gaussian-type wave function [25]

$$\begin{aligned}\varphi' &= \varphi'(x_2, p'_\perp) = 4 \left(\frac{\pi}{\beta'^2} \right)^{\frac{3}{4}} \sqrt{\frac{dp'_z}{dx_2}} \exp \left(-\frac{p_z'^2 + p_\perp'^2}{2\beta'^2} \right), \\ \varphi'_p &= \varphi'_p(x_2, p'_\perp) = \sqrt{\frac{2}{\beta'^2}} \varphi', \quad \frac{dp'_z}{dx_2} = \frac{e'_1 e_2}{x_1 x_2 M'_0}.\end{aligned}\tag{29}$$

The input parameters m_q and β in the Gaussian-type wave function (29) are shown in Table I. The constituent quark masses are close to those used in the literature [14, 15, 17, 18, 19, 20]. The parameter β' , which describes the momentum distribution, is expected to be of order Λ_{QCD} . These parameters β 's are fixed by the decay constants whose analytic expressions in the covariant light-front model are given in [18]. The decay constant $f_{J/\psi}$ can be determined by the leptonic decay width:

$$\Gamma_{ee} \equiv \Gamma(J/\psi \rightarrow e^+ e^-) = \frac{4\pi\alpha_{em}^2 Q_c^2 f_{J/\psi}^2}{3m_{J/\psi}},\tag{30}$$

where $Q_c = 2/3$, denotes the electric charge of the charm quark. Using the measured results for the electronic width of J/ψ [26]:

$$\Gamma_{ee} = (5.55 \pm 0.14 \pm 0.02) \text{ keV},\tag{31}$$

we obtain $f_{J/\psi} = 0.416 \pm 0.05 \text{ GeV}$. As for the decay constant for B_c and X , we use $f_{B_c} = 0.398^{+0.054}_{-0.055} \text{ GeV}$, and $|f_{X(3872)}| = 0.329^{+0.111}_{-0.095} \text{ GeV}$. For the light pseudoscalars, we use $f_\pi = 0.132 \text{ GeV}$ and $f_K = 0.16 \text{ GeV}$. The determined results for β are listed in Table I.

In the factorization approach, the decay amplitude is expressed as a product of the short distance Wilson coefficients, the form factors and meson decay constant. The latter two are physical values which are scale-independent. But the Wilson coefficient a_1 of the four quark operators depends on the factorization scale. This directly leads to the scale dependence of

the decay amplitude. But as we have shown in Ref. [27], the numerical value of a_1 is not very sensitive to the scale thus we use $a_1 = 1.1$ in this work. The CKM matrix elements, lifetime of B_c meson and the masses of the hadrons are chosen from the Particle Data Group [26] as:

$$|V_{cb}| = 0.0416 \pm 0.0006, \quad |V_{ud}| = 0.97377 \pm 0.00027, \quad |V_{us}| = 0.2257 \pm 0.021, \quad (32)$$

$$m_{B_c} = 6.286 \text{ GeV}, \tau_{B_c} = (0.46^{+0.18}_{-0.16}) \times 10^{-12} \text{ s}, \quad (33)$$

$$m_{J/\psi} = 3.097 \text{ GeV}, \quad m_X = 3.872 \text{ GeV}. \quad (34)$$

The uncertainties in the above CKM matrix elements are small, thus induce small errors to the decay width and we will neglect these uncertainties.

Using the above inputs, we can calculate the form factors directly. As in Refs. [17, 18], because of the condition $q^+ = 0$ we have imposed during the course of calculation, form factors are known only for spacelike momentum transfer $q^2 = -q_\perp^2 \leq 0$. But only the timelike form factors are relevant for the physical decay processes. It has been proposed in [14] to recast the form factors as explicit functions of q^2 in the spacelike region and then analytically extrapolate them to the timelike region. In the exclusive non-leptonic decays, only the form factor at maximally recoiling ($q^2 \simeq 0$) is required therefore we do not need to discuss the dependence on the momentum transfer here. After calculation, the results for the $B_c \rightarrow J/\psi$ and $B_c \rightarrow X$ (assuming X as a 1^{++} state) form factors are

$$\begin{aligned} V^{PV}(0) &= 0.87^{+0.00+0.01}_{-0.02-0.00}, & A_0^{PV}(0) &= A_3^{PV}(0) = 0.57^{+0.01+0.00}_{-0.02-0.00}, \\ A_1^{PV}(0) &= 0.55^{+0.01+0.00}_{-0.03-0.00}, & A_2^{PV}(0) &= 0.51^{+0.03+0.00}_{-0.04-0.00}, \\ A^{PA}(0) &= 0.36^{+0.02+0.01}_{-0.02-0.03}, & V_0^{PA}(0) &= V_3^{PA}(0) = 0.18^{+0.01+0.01}_{-0.02-0.02}, \\ V_1^{PA}(0) &= 1.15^{+0.03+0.03}_{-0.04-0.06}, & V_2^{PA}(0) &= 0.13^{+0.02+0.00}_{-0.02-0.01}, \end{aligned} \quad (35)$$

where the first uncertainty is from the decay constant of the B_c meson and the latter is from the charmonium decay constant. In Table II, we make a comparison of our results with the previous studies. We can see that our results are slightly smaller than the results from the three points sum rule and quark model, but the light-cone sum rule predictions are quite different from the others.

Using the results for $B_c \rightarrow J/\psi$ form factors, we obtain the branching ratio of $B_c^- \rightarrow J/\psi \pi^- (K^-)$:

$$\begin{aligned} \mathcal{BR}(B_c^- \rightarrow J/\psi \pi^-) &= (2.0^{+0.8+0.0+0.0}_{-0.7-0.1-0.0}) \times 10^{-3}, \\ \mathcal{BR}(B_c^- \rightarrow J/\psi K^-) &= (1.6^{+0.6+0.0+0.0}_{-0.6-0.1-0.0}) \times 10^{-4}, \end{aligned} \quad (36)$$

where the uncertainties are from the large errors in the lifetime of B_c meson, the decay constant of B_c meson and the charmonium decay constants. In the literature, these decays have received extensively study [31] and the range of the branching ratios is

$$\begin{aligned} \mathcal{BR}(B_c^- \rightarrow J/\psi \pi^-) &= (0.06 - 0.18)\%, \\ \mathcal{BR}(B_c^- \rightarrow J/\psi K^-) &= (0.005 - 0.014)\%, \end{aligned} \quad (37)$$

TABLE II: The values of the form factors of $B_c \rightarrow J/\psi$ at $q^2 = 0$ in comparison with the estimates in the three points sum rule (3PSR) (with the Coloumb corrections included) [28], in the quark model (QM) [29] and the light-cone sum rules (LCSR) [30]

	A_1	A_2	V
3PSR [28]	0.63	0.69	1.03
QM [29]	0.68	0.66	0.96
LCSR [30]	0.75	1.69	1.69
This work	$0.55^{+0.01+0.00}_{-0.03-0.00}$	$0.51^{+0.03+0.00}_{-0.04-0.00}$	$0.87^{+0.00+0.01}_{-0.02-0.00}$

which are consistent with ours. Assuming X as a 1^{++} state, the branching ratios of $B_c^- \rightarrow X(3872)\pi^-(K^-)$ are

$$\begin{aligned}\mathcal{BR}(B_c^- \rightarrow X(3872)\pi^-) &= (1.7^{+0.7+0.1+0.4}_{-0.6-0.2-0.4}) \times 10^{-4}, \\ \mathcal{BR}(B_c^- \rightarrow X(3872)K^-) &= (1.3^{+0.5+0.1+0.3}_{-0.5-0.2-0.3}) \times 10^{-5}.\end{aligned}\quad (38)$$

These results are one order magnitude smaller than the branching ratio of $B_c^- \rightarrow J/\psi\pi^-$ and $B_c^- \rightarrow J/\psi K^-$, respectively. From the decay width formulae in Eq. (28), we know that the $B_c \rightarrow J/\psi P$ branching ratios are proportional to the form factor $|A^{PV}(0)|^2$ while for $B_c \rightarrow X(1^{++})P$, the decay width are proportional to $|V^{PA}(0)|^2$. The $B_c \rightarrow X$ form factor $V^{PA}(0)$ is only 1/3 of $|A^{PV}(0)|$ shown in Eq. (35), which induces the one order magnitude difference for these two kinds of decays.

For $B_c \rightarrow VV$ decays, there are three different polarizations. According to the power counting rule in the standard model [32], the longitudinal polarization dominates in the decay processes, while other polarizations suffer one or two orders of Λ_{QCD}/m_B or m_c/m_B suppressions which arise from the quark helicity flip. It is found that the annihilation diagrams with the operator O_6 could violate this power counting rule [33]. However in the $B_c^- \rightarrow J/\psi\rho^-(K^{*-})$ decays, there are only emission type contributions thus the power counting rule should work well. If we neglect the $m_\rho^2/m_{B_c}^2$ terms in the polarization vector, the formulae for the branching ratios of $B \rightarrow VV$ is the same as eq. (28) with the replacement of the decay constant: $f_P \rightarrow f_V$. Using $f_\rho = 0.209$ GeV and $f_{K^*} = 0.217$ GeV, we obtain the corresponding branching ratios as:

$$\begin{aligned}\mathcal{BR}(B_c^- \rightarrow J/\psi\rho^-) &= (5.0^{+2.0+0.1+0.1}_{-1.7-0.0-0.0}) \times 10^{-3}, \\ \mathcal{BR}(B_c^- \rightarrow J/\psi K^{*-}) &= (2.9^{+1.1+0.0+0.0}_{-1.0-0.0-0.0}) \times 10^{-4}, \\ \mathcal{BR}(B_c^- \rightarrow X(3872)\rho^-) &= (4.1^{+1.6+0.3+0.1}_{-1.4-0.1-0.1}) \times 10^{-4}, \\ \mathcal{BR}(B_c^- \rightarrow X(3872)K^{*-}) &= (2.4^{+0.9+0.2+0.5}_{-0.8-0.3-0.5}) \times 10^{-5}.\end{aligned}\quad (39)$$

Our above calculation is based on the 1^{++} charmonium description for X . The charmonium states with other quantum numbers can also be studied similarly in this approach.

If the quantum numbers of $X(3872)$ are changed to 1^{--} , the large form factor $A_0^{B_c \rightarrow X}$ can enhance the production rates dramatically as:

$$\begin{aligned}
\mathcal{BR}(B_c^- \rightarrow X(3872)\pi^-) &= (1.4_{-0.5-0.0-0.5}^{+0.6+0.0+0.4}) \times 10^{-3}, \\
\mathcal{BR}(B_c^- \rightarrow X(3872)K^-) &= (1.1_{-0.4-0.0-0.4}^{+0.4+0.0+0.3}) \times 10^{-4}, \\
\mathcal{BR}(B_c^- \rightarrow X(3872)\rho^-) &= (3.5_{-1.2-0.2-1.2}^{+1.4+0.0+1.1}) \times 10^{-3}, \\
\mathcal{BR}(B_c^- \rightarrow X(3872)K^{*-}) &= (2.0_{-0.7-0.1-0.7}^{+0.8+0.0+0.6}) \times 10^{-4}.
\end{aligned} \tag{40}$$

Comparing the above equations with eqs. (38) and (39), we can see that the production rates are enhanced by about one order magnitude. The large branching ratios and large difference between different quantum number of state X can be easily used at LHCb experiment to test the charmonium description for X .

IV. CONCLUSION

In the covariant light-front quark model, we study the form factors of $B_c \rightarrow J/\psi$ and $B_c \rightarrow X$ transitions at maximum recoiling. The factorization of exclusive processes $B_c^- \rightarrow J/\psi\pi^-(K^-)$ and $B_c^- \rightarrow X(3872)\pi^-(K^-)$ can be proved to all orders of the strong coupling constant just as the proof in $\bar{B}^0 \rightarrow D^+\pi^-$ and $B^- \rightarrow D^0\pi^-$. Therefore the decay width of these decays can be simply calculated in the naive factorization approach utilizing the form factors. Our results for the branching ratio of $B_c \rightarrow J/\psi\pi^-(K^-)$ are consistent with the previous studies considering the uncertainties. The study of these exclusive processes may greatly improve our understanding on the B_c meson exclusive hadronic decays, and the corresponding theory describing them as well. In our calculation, identifying $X(3872)$ as a 1^{++} charmonium state, we obtain $\mathcal{BR}(B_c^- \rightarrow X(3872)\pi^-) = (1.7_{-0.6-0.2-0.4}^{+0.7+0.1+0.4}) \times 10^{-4}$ and $\mathcal{BR}(B_c^- \rightarrow X(3872)K^-) = (1.3_{-0.5-0.2-0.3}^{+0.5+0.1+0.3}) \times 10^{-5}$. If assuming X as a 1^{--} state, the branching ratios are one order magnitude larger. This large difference can be easily used by the LHCb experiment to test the different charmonium description for X .

Acknowledgement

We thank H.Y. Cheng, C.K. Chua and Y.M. Wang for useful discussions. This work is partly supported by National Science Foundation of China under Grant No. 10475085 and 10625525.

-
- [1] S. K. Choi *et al.* [Belle Collaboration], Phys. Rev. Lett. **91**, 262001 (2003).
 - [2] D. Acosta *et al.* [CDF II Collaboration], Phys. Rev. Lett. **93**, 072001 (2004);
V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **93**, 162002 (2004);

- B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **71**, 071103 (2005).
- [3] K. Abe *et al.*, arXiv:hep-ex/0505038.
- [4] T. Barnes and S. Godfrey, Phys. Rev. D **69**, 054008(2004).
- [5] E.S. Swanson, Phys. Rept. **429**, 243(2006).
- [6] B.A. Li, Phys. Lett. B **605**, 306 (2005).
- [7] K.K. Seth, Phys. Lett. B **612**, 1 (2005).
- [8] L. Maiani, F. Piccinini, A.D. Polosa, and V. Riquer, Phys. Rev. D **71**, 014028 (2005);
T.W. Chiu and T.H. Hsieh [TWQCD Collaboration], arXiv:hep-ph/0603207.
- [9] N.A. Törnqvist, arXiv:hep-ph/0308277;
F.E. Close, and P.R. Page, Phys. Lett. B **578**, 119 (2004);
S. Pakvasa and M. Suzuki, Phys. Lett. B **579**, 67 (2004);
E.S. Swanson, Phys. Lett. B **588**, 189 (2004).
- [10] Y.Chen *et al.* [CLQCD Collaboration], arXiv: hep-ph/0701021.
- [11] H. Y. Cheng, Y. Y. Keum and K. C. Yang, Phys. Rev. D **65**, 094023 (2002).
- [12] C.W. Bauer, D. Pirjol and I.W. Stewart, Phys. Rev. Lett. **87**, 201806(2001).
- [13] S.J. Brodsky, H.C. Pauli and S.S. Pinsky, Phys. Rep. **301**, 299(1998).
- [14] W. Jaus, Phys. Rev. D **41**, 3394(1990); Phys. Rev. D **44**, 2851(1991).
- [15] H.-Y. Cheng, C.-Y. Cheung and C.-W. Hwang, Phys. Rev. D **55** 1559 (1997).
- [16] H.-M. Choi, C.-R. Ji, and L.S. Kisslinger, Phys. Rev. D **65**, 074032(2002).
- [17] W. Jaus, Phys. Rev. D **60**, 054026(1999).
- [18] H.-Y. Cheng, C.-K. Chua and C.-W. Hwang, Phys. Rev. D **69**, 074025, (2004).
- [19] H. Y. Cheng and C. K. Chua, Phys. Rev. D **69**, 094007 (2004).
- [20] C. W. Hwang and Z. T. Wei, J. Phys. G **34**, 687 (2007).
- [21] For a review, see G. Buchalla, A.J. Buras, M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
- [22] C.W. Bauer, S. Fleming, D. Pirjol and I.W. Stewart, Phys. Rev. D **63**, 114020(2001).
- [23] M. Beneke, A.P. Chapovsky, M. Diehl and Th. Feldmann, Nucl. Phys. B **643**, 431(2002).
- [24] M. Wirbel, B. Stech, M. Bauer Z. Phys. C **29**, 637(1985).
- [25] P. L. Chung, F. Coester, and W. N. Polyzou, Phys. Lett. B **205**, 545 (1988).
- [26] Particle Data Group, W.-M. Yao *et al.*, J. Phys. G **33**,1 (2006).
- [27] A. Ali *et al.*, arXiv: hep-ph/0703162.
- [28] V. V. Kiselev, A.K. Likhoded, and A. I. Onishchenko, Nucl. Phys. B **569**, 473 (2000);
V. V. Kiselev, A. E. Kovalsky, A. K. Likhoded, Nucl. Phys. B **585**, 353 (2000).
- [29] M. A. Ivanov, J. G. Korner, P. Santorelli, Phys. Rev. D **63**, 074010 (2001).
- [30] T. Huang and F. Zuo, arXiv: hep-ph/0702147.
- [31] C.H. Chang, Y.Q. Chen, Phys. Rev. D **49**, 3399(1994); C.H. Chang, Y.Q. Chen, G.L. Wang,
and H.S. Zong, Commun. Theor. Phys. **35**, 395(2001); Phys. Rev. D **65**, 014017(2002);
A.Y. Anisimov, P.Y. Kulikov, I.M. Narodetsky, and K. A. Ter-Martirosian, Yad. Fiz. **62**,

- 1868(1999) [Phys. Atom. Nucl. **62**, 1739(1999)];
P. Colangelo, and F. De Fazio, Phys. Rev. D**61**, 034012(2000);
A. Abd El-Hady, J.H. Munoz, and J.P. Vary, Phys. Rev. D**62**, 014019(2000);
G. Lopez Castro, H.B. Mayorga, and J.H. Muñoz, J. Phys. G**28**, 2241(2002);
V.V. Kiselev, hep-ph/0211021;
V.V. Kiselev, O.N. Pakhomova, and V.A. Saleev, J. Phys. G **28**, 595(2002);
D. Ebert, R.N. Faustov, and V.O. Galkin, Phys. Rev. D**68**, 094020(2003);
M.A. Ivanov, J.G. Körner, and P. Santorelli, Phys. Rev. D**73**, 054024(2006);
E. Hernández, J. Nieves and J. M. Verde-Velasco, Phys. Rev. D**74**, 074008 (2006).
[32] A.L. Kagan, Phys. Lett. B**601**,151 (2004); H.-n. Li and S. Mishima, Phys. Rev. D**71**, 054025 (2005); H.-n. Li, Phys. Lett. B**622**, 63 (2005).
[33] Y.Y. Keum, H.-n. Li, and A.I. Sanda, Phys. Lett. B**504**, 6 (2001); Phys. Rev. D **63**, 054008 (2001);
C.-D. Lu, K. Ukai and M.-Z. Yang, Phys. Rev. D**63**, 074009 (2001).